

**Anti-Trypsin (human)
Mouse monoclonal antibody**

Subclass: IgG1/k

PRODUCT NO.

HYB 021-03

Clone: 7C8

PRESENTATION

Preparation: Protein-A/G purified

Content: Available in 200 μ L and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.

Solvent: 0.01 M phosphate buffer, pH 7.4, containing 0.5 M NaCl and 15 mM sodium azide

Storage: 4-8°C without exposure to light. No precautions necessary during handling.

ANTIGEN

Trypsin is a 24 kDa enzyme and is a member of the serine proteinase family. It is produced in the pancreas as an inactive precursor trypsinogen. The active enzyme is located in the gastrointestinal tract where it degrades proteins to large peptides. Trypsin predominantly cleaves proteins at the carboxyl side of the amino acids lysine and arginine. A common trypsin substrate is chymotrypsinogen which is cleaved to produce active chymotrypsin.

High levels of immunoreactive trypsin in the bloodstream can indicate pancreatic malfunction and this indicator is used to screen for and diagnose Cystic Fibrosis (1).

IMMUNOGEN

Trypsin purified from human pancreatic tissue and adsorbed onto aluminum hydroxide gel (2)

SPECIFICITY

HYB 021-03 is specific for human trypsin, trypsinogen and trypsin in complex with α_1 -antitrypsin. HYB 021-03 cross-reacts weakly with bovine trypsin.

EPI TOPE SPECIFICITY

Epitope differs from that of HYB 021-08

REACTIVITY

HYB 021-03 reacts with human trypsin coated directly onto a microtiter plate and can be used as both a catching and a detection antibody in sandwich ELISA.

HYB 021-03 recognizes human trypsin in Western blotting under reducing conditions.

CULTURE MEDIUM

RPMI 1640 with 10% fetal calf serum

FUSION PARTNER

X63-Ag8.653

IMMUNIZATION

Female CF1 x BALB/c mice immunized by intraperitoneal injection

APPLICATION

Method	Usability	References
ELISA	Yes	
Immunoblotting	Yes	
Immunohistochemistry	Not determined	

REFERENCES

1. Crossley JR, Smith PA, Edgar BW, Gluckman PD, Elliott RB (1981) Neonatal screening for cystic fibrosis, using immunoreactive trypsin assay in dried blood spots. Clin Chim Acta 113:111-121

2. Feinstein G, Hofstein R, Koifmann J, Sokolovsky M (1974) Human pancreatic proteolytic enzymes and protein inhibitors. Isolation and molecular properties. Eur J Biochem 43:569-581.

CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its products.