

**Anti-Gc-globulin (human)  
Mouse monoclonal antibody**

 Subclass: IgG<sub>1</sub>/k

PRODUCT NO.

**HYB 249-10**

Clone: 4B9

PRESENTATION

Preparation: Protein-A/G purified

Content: Available in 200 µL and 1 mL volumes, 1 mg/mL

Solvent: 0.01 M phosphate buffer, pH 7.4, containing 0.5 M NaCl and 15 mM sodium azide

Storage: In the dark at 4-8°C

ANTIGEN

Gc-globulin is a plasma protein produced in the liver. Amongst its ligands are vitamin D, thus Gc-globulin is also called vitamin D-binding protein. Gc-globulin is furthermore part of the actin scavenging system, binding and removing monomeric actin from the blood stream. The molecular mass of Gc-globulin is approximately 50 kDa. The concentration of Gc-globulin in human plasma is app. 400 µg/ml (1).

IMMUNOGEN

Gc-globulin isolated from human plasma adsorbed onto aluminum hydroxide gel

SPECIFICITY

HYB 249-10 is specific for human Gc-globulin

EPI TOPE SPECIFICITY

Epitope specificity differs from that of HYB 249-01, HYB 249-02 and HYB 249-05.

REACTIVITY

HYB 249-10 reacts strongly with Gc-globulin. Strong reaction is seen in ELISA with Gc-globulin coated directly onto the microtiter plate, and also when tested in sandwich ELISA in combination with a polyclonal antibody against Gc-globulin (e.g. DAKO A 0021). In Western blotting after SDS-PAGE, HYB 249-10 reacts with Gc-globulin in reduced as well as in unreduced forms.

CULTURE MEDIUM

RPMI 1640 with 10% fetal calf serum

FUSION PARTNER

X63-Ag8.653

IMMUNIZATION

Female CF1 x BALB/c mice immunized by intraperitoneal injection

APPLICATION

Method	Usability	Dilution guideline	References
ELISA	Yes	1/20,000	
Immunoblotting	Yes		
Immunohistochemistry	Not determined		

The dilution guideline for ELISA is based on sandwich ELISA in combination with a polyclonal antibody against the antigen. Users should determine the optimal dilutions for their own purpose.

REFERENCES

1. Masuda S, Okano T, Osawa K, Shinjo M, Suematsu T, Kobayashi T (1989) Concentrations of vitamin D-binding protein and vitamin D metabolites in plasma of patients with liver cirrhosis. J Nutr Sci Vitaminol 35:225-34.

**CONDITIONS**

All products are supplied on the understanding that they are for in vitro use only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The animals from which this product was derived have not been exposed to or inoculated with any livestock or poultry disease agents exotic to the United States or Western Europe, and did not originate from facilities where work with exotic disease agents affecting livestock or avian species is carried out.