

**Anti Complement component C2 (human)
mouse monoclonal antibody**Subclass: IgG₁/k

PRODUCT NO.	HYB 050-05
PRESENTATION	Preparation: Protein-A/G purified Content: Available in 200 µL and 1 mL, 1 mg/mL Solvent: 0.01 M phosphate buffer, pH 7.4, with 0.5 M NaCl and 15 mM sodium azide Storage: In the dark at 4-8°C
ANTIGEN	Complement component C2 is a component of the classical complement pathway and is involved in activation of C3 and C5. C2 consists of a single polypeptide chain with a molecular mass of 102 kDa (2).
IMMUNOGEN	Complement component C2 isolated from human plasma (1)
SPECIFICITY	HYB 050-05 is specific for C2 from human plasma/serum. Crossreactivity to C2 from other species has not been tested.
EPI TOPE SPECIFICITY	Epitope specificity differs from that of HYB 050-04 and HYB 050-08 as determined by inhibition ELISA.
REACTIVITY	HYB 050-05 reacts strongly with C2. Strong reaction is seen with human serum/plasma and when used as detecting antibody in sandwich ELISA in combination with a polyclonal antibody against C2. In Western blotting after SDS-PAGE, HYB 050-05 reacts with C2 and a subfraction of C2 believed to be C2b, but only under non-reducing conditions. HYB 050-05 is suitable for Sepharose column purification of C2 from plasma (3,4).
CULTURE MEDIUM	RPMI 1640 with 10% fetal calf serum
FUSION PARTNER	X63-Ag8.653.
IMMUNIZATION	Female CF1 x BALB/c mice immunized i.p. with immunogen adsorbed onto Al(OH) ₃

APPLICATION

Method	Usability	Dilution guideline	References
ELISA	Yes	1:4000	1
Immunoblotting	Yes	1:500	1
Immunohistochemistry	Not determined		

The dilution guideline for ELISA is based on sandwich ELISA in combination with a polyclonal antibody against the antigen. Users should determine the optimal dilutions for their own purpose.

REFERENCES

1. Stenbaek EI, Koch C, Barkholt V, Welinder KG (1986) Human complement component C2: production and characterization of polyclonal and monoclonal antibodies against C2. *Mol Immunol* 23:879-886.
2. Law SKA, Reid KBM (1988) Complement. In: *In Focus* (Ed. Male D) IRL Press: Oxford.
3. Laich A, Moffatt B, Wong KHN, Hickling TP, Koch C, Sim RB (2001) Purification of second component of human complement, C2, by antibody affinity chromatography. *Intern J Bio-Chromatography* 6:151-162.
4. Laich A, Sim RB (2001) Complement C4bC2 complex formation: an investigation by surface plasmon resonance. *Biochim Biophys Acta* 1544:96-112.

CONDITIONS

All products are supplied on the understanding that they are for in vitro use only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The animals from which this product was derived have not been exposed to or inoculated with any livestock or poultry disease agents exotic to the United States or Western Europe, and did not originate from facilities where work with exotic disease agents affecting livestock or avian species is carried out.