



**Anti Complement component C5 (human)
Mouse Monoclonal Antibody**

Subclass: IgG₁/κ

PRODUCT NO.	HYB 029-02																
PRESENTATION	Preparation: Protein-A purified Content: 1 ml, 1 mg/mL Solvent: 0.01 M phosphate buffer, pH 7.4, with 0.5 M NaCl and 15mM sodium azide Storage: In the dark at 4-8°C																
ANTIGEN	C5 is part of the membrane attack complex (MAC) of the complement system. C5 is split into C5a og C5b by the C5 convertases from the classical and the alternative complement pathway and the C5b fragment initiates the assembly of the terminal MAC (C5b-C9) (1). The molecular mass of C5 is 190 kDa.																
IMMUNOGEN	C5 isolated from human plasma																
SPECIFICITY	HYB 029-02 is specific for human C5. No reaction is seen with plasma from patients deficient in C5.																
EPITOPE SPECIFICITY	Epitope specificity differs from that of HYB 029-03 but slightly overlap as determined by inhibition ELISA.																
REACTIVITY	HYB 029-02 reacts strongly with C5. Strong reaction is seen with normal human plasma, and no reaction is seen with plasma from patients deficient in C5, when tested in sandwich ELISA in combination with a polyclonal antibody against C5 (eg. DAKO A 055). In Western blotting after SDS-PAGE, HYB 029-02 reacts with C5 in normal human plasma (PEG precipitated C5) in nonreduced form only and no reaction is seen with plasma from C5 deficient patients.																
CULTURE MEDIUM	Dulbecco's modified Eagle's medium with 10% fetal calf serum																
FUSION PARTNER	X63-Ag8.653.																
IMMUNIZATION	Female CF1 x Balb/c mice, immunized intraperitoneally with antigen adsorbed onto Al(OH) ₃ .																
APPLICATION	<table border="1"> <thead> <tr> <th>Method</th> <th>Usability</th> <th>Dilution guideline</th> <th>References</th> </tr> </thead> <tbody> <tr> <td>ELISA</td> <td>Yes</td> <td>1:6000</td> <td></td> </tr> <tr> <td>Immunoblotting</td> <td>Yes</td> <td></td> <td></td> </tr> <tr> <td>Immunohistochemistry</td> <td>Not determined</td> <td></td> <td></td> </tr> </tbody> </table> <p>The dilution guideline for ELISA is based on sandwich ELISA in combination with a polyclonal antibody against the antigen. Users should determine the optimal dilutions for their own purpose.</p>	Method	Usability	Dilution guideline	References	ELISA	Yes	1:6000		Immunoblotting	Yes			Immunohistochemistry	Not determined		
Method	Usability	Dilution guideline	References														
ELISA	Yes	1:6000															
Immunoblotting	Yes																
Immunohistochemistry	Not determined																
REFERENCES	1. Law SKA, Reid KBM (1988) Complement. In: In Focus (Ed. Male D) IRL Press: Oxford.																

CONDITIONS

All products are supplied on the understanding that they are for in vitro use only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The animals from which this product was derived have not been exposed to or inoculated with any livestock or poultry disease agents exotic to the United States or Western Europe, and did not originate from facilities where work with exotic disease agents affecting livestock or avian species is carried out.