

**Anti Complement component C3d (human)  
Mouse Monoclonal Antibody**Subclass: IgG<sub>1</sub>/k

PRODUCT NO.	<b>HAV 003-05</b>
PRESENTATION	Preparation: Protein-A/G purified Content: 1 ml, 1 mg/mL Solvent: 0.01 M phosphate buffer, pH 7.4, with 0.5 M NaCl and 15mM sodium azide Storage: In the dark at 4-8°C
ANTIGEN	C3 is a key component of the complement system since classical and alternative activation pathways merge at the C3 activation step when C3 is split into C3a and C3b. The molecular mass of C3 is 185 kDa and it consists of two chains (110 kDa and 75 kDa) held together by disulfide bonds (1,2).
IMMUNOGEN	C3 isolated from human plasma
SPECIFICITY	HAV 003-05 is specific for human C3d, a fragment of the alpha chain (3)
EPITOPE SPECIFICITY	Epitope specificity differs from that of HAV 004-01 and HYB 030-08
REACTIVITY	HAV 003-05 reacts strongly with C3. Strong reaction is seen in ELISA with human C3 coated directly onto the microtiter well, or when used as detection antibody in sandwich ELISA in combination with a polyclonal C3 antibody (e.g. DAKO A 062). HAV 003-05 also reacts with C3b deposited on coated antibody molecules. In Western blotting after SDS-PAGE, HAV 003-05 reacts with C3 in both reduced and unreduced forms.
CULTURE MEDIUM	RPMI 1640 with 10% fetal calf serum
FUSION PARTNER	X63-Ag8.653.

IMMUNIZATION Female CF1 x BALB/c mice immunized i.p. with immunogen adsorbed onto Al(OH)<sub>3</sub>

Method	Usability	Dilution guideline	References
ELISA	Yes	1:16,000	4
Immunoblotting	Yes		
Immunohistochemistry	Not determined		

The dilution guideline for ELISA is based on sandwich ELISA in combination with a polyclonal antibody against the antigen. Users should determine the optimal dilutions for their own purpose.

## REFERENCES

1. Law SKA, Reid KBM (1988) Complement. In: In Focus (Ed. Male D) IRL Press: Oxford.
2. Morley BJ and Walport MJ (2000) The Complement FactsBook. Academic Press, London, UK.
3. Nagar B, Jones RG, Diefenbach RJ, Isenman DE, Rini JM (1998) X-ray crystal structure of C3d: a C3 fragment and ligand for complement receptor 2. Science 280:1277-1281.
4. Koch C, Behrendt N (1986) A novel polymorphism of human complement component C3 detected by means of a monoclonal antibody. Immunogenetics 23:322-325.

## CONDITIONS

All products are supplied on the understanding that they are for in vitro use only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The animals from which this product was derived have not been exposed to or inoculated with any livestock or poultry disease agents exotic to the United States or Western Europe, and did not originate from facilities where work with exotic disease agents affecting livestock or avian species is carried out.